

27.05.03 #2



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

REC'D 06 JUN 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02356081.6

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk



Anmeldung Nr:  
Application no.: 02356081.6  
Demande no:

Anmeldetag:  
Date of filing: 30.04.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

AVENTIS ANIMAL NUTRITION S.A.  
42 Avenue Aristide Briand  
92160 Antony  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Animal feed supplement

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A23K/

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

## ANIMAL FEED SUPPLEMENT

The present invention relates to a novel animal feed supplement.

5 Methionine is a well known additive for animal feed, in particular for ruminants. Methionine may be added directly to the animal feed. The free form of this compound however is degraded rapidly by bacteria in the rumen and consequently only a small portion of the original methionine is able to enter the bloodstream. An analogue of methionine is also often used, namely the hydroxy analogue having the chemical name 2-hydroxy-4-(methylthio) butanoic acid,  
10 generally referred to as HMB. This product is used in the liquid form and is also largely degraded in the rumen.

A commercially known product sold as Smartamine sold by Aventis Animal Nutrition is effective in providing methionine to the animal and tests have shown that the methionine enters the blood stream in 10 hours after  
15 administration. Recent research has also shown that ceratin esters of methionine or HMB may also be successfully used to provide the necessary bio-available methionine to ruminants. WO 00/28835 discloses the use of the ester form of the hydroxy analogue and in particular this document teaches that at least 50% of bio-available methionine may be provided through the administration of the iso  
20 propyl ester of the hydroxy analogue of methionine.

We have now found that when the hydroxy methionine analogue, and in particular the iso propyl ester, is provided in the solid form and admixed with the animal feed, the product shows good bioavailable results at least equivalent to the liquid form.

25 Accordingly, the present invention provides an animal feed supplement in the solid form said feed supplement comprising 2-hydroxy-4-(methylthio) butanoic acid or an ester thereof and a porous carrier material wherein the 2-hydroxy-4-(methylthio) butanoic acid or ester thereof accounts for at least 20% by weight of the total composition of the feed supplement.

The present invention provides the advantage over the previously known animal feed supplements in that the use of the particular feed supplement not only provides a large amount of methionine to the animal but that methionine is found to be present in the bloodstream after a short period of time when compared to the commercially available methionine sources.

The present invention provides an animal feed supplement that is in the solid form. For the purposes of the present invention solid form is defined as powder wherein the particles or granules of powder have a size range of from 200 to 400 microns.

The feed supplement of the present invention comprise 2-hydroxy-4-(methylthio) butanoic acid or an ester thereof. Suitable esters include linear or branched alkyl esters such as methyl ester, ethyl ester, n-propyl ester, isopropyl ester, butyl esters, namely n-butyl ester, sec butyl ester, isobutyl ester and tertiary butyl ester, pentyl esters and hexyl esters, especially n-pentyl, isopentyl, n-hexyl and isohexyl esters.

Preferably, the ester is a branched alkyl ester, especially the isopropyl ester and the tertiary butyl ester. In particular, it has been found that a solid feed supplement comprising the isopropyl ester of the hydroxy analogue of methioinine displays a bio-availability of methionine of more than 50%.

The 2-hydroxy-4-(methylthio) butanoic acid or the ester thereof accounts for at least 20% by weight of the total feed composition. Suitably, this compound is present in an amount of from 20 to 70% by weight, preferably from 25 to 50% by weight.

The feed supplement of the present invention comprise a porous carrier material. Suitably the carrier material has a porosity of at least 0.25 ml/g, preferably from 0.4 to 2.0 ml/g. Suitable carrier materials include clays such as vermiculite, sepiolite and perlite clays; silicates and silica such as magnesium silicates and zeolites. The preferred material is a clay, especially sepiolite, commercially available as Exal-H.

The granule may comprise additional components such as an anti-caking agent, an anti-oxidising agent and a surfactant. Anti-caking agents suitable for use in the present invention include silicas, talcum, calcium phosphate, calcium carbonate and magnesium stearate. The preferred anti-caking agent is a compound having a high surface area, for example at least 100 m<sup>2</sup>/g. Preferably, the anti-caking agent is silica. The anti-caking agent is suitable present in an amount of from 0 to 2% by weight.

As regards the anti-oxidising agent, suitable agents include tocopherol, ascorbic acid, ethoxyquine, propyl, octyl or dodecyl gallate, butyl hydroxy anisole (commonly referred to as BHA) and butylated hydroxytoluene (commonly referred to as BHT). The anti-oxidant may be present in the solid in an amount of from 0 to 3% by weight of the granule.

Surfactants suitable for use in the present invention include sulphates such as sodium alkyl sulphate or lauryl sulphate, polysorbates such as polyoxy ethylene sorbitol fatty acid esters. The surfactant is present in an amount of from 0 to 5% by weight.

The granule may contain water although it is preferred that the granule is essentially dry. The water content may be from 0 to 10 % by weight.

A particularly preferred granule of the present invention comprises 25 to 30% by weight of the isopropyl ester of the hydroxy analogue of methionine, and 75 to 70% of sepiolite.

The solid particle may suitably have an average medium diameter of from 200 to 400 microns, preferably from 275 to 325 microns and a bulk density suitably of from 700 to 950 kg/m<sup>3</sup>, preferably from 800 to 900 kg/m<sup>3</sup>.

The feed supplement of the present invention may be prepared by impregnation of the carrier material with the liquid methionine containing product optionally followed by addition of the anti-caking agent. This procedure may be carried out by mixing the components using any suitable blender or mixer. The mixing may be carried out at a temperature of from 20 to 80°C, preferably from

40 to 70 °C. The speed of mixing will be determined by the desired size of the resulting powder. Alternatively, the mixing may be carried out in a fluidised bed.

The feed supplement of the present invention may be incorporated into the food of the animal and according to another aspect of the present invention there is provided an animal feed comprising a feed supplement as herein before defined.

The present invention will now be described in detail with reference to the following examples:

In the following preparative work, the following standard tests were carried out on the solid supplement :

1. Particle Size: The size of the solid particle was determined by passing 100g of the granules through Retsch sieves with 1.5 mm of amplitude during a period of 10 minutes.

2. Bulk and Tapped Density: The bulk and tapped density were determined by powder volumenometer (230ml).

3. Carr Index - This index is defined as :

$$\frac{(\text{Bulk Density} - \text{Tapped Density})}{\text{Bulk Density}}$$

4. Flow Index: The flow Index was determined by a flow tester which is the ring shear stress of Schulze (device reference RST-01.01 Dr Ing. Dietmar Schulze, Wolfenbuttel, Germany). Jenike Index is defined by the ratio of principal stress at a steady flow to the unconfined yield strength.

#### Example 1: Preparation of Feed Supplement Containing 2-hydroxy-4-(methylthio) butanoic acid

A weight of 35kg of sepiolite clay (70 weight %) was placed in a 130 litres Lodige plow mixer. The clay was stirred for five minutes. 15kg of liquid 2-hydroxy-4-(methylthio) butanoic acid (30weight %) was heated to 60°C and then poured into the mixer at a flow rate of 1000kg/hour. The resulting mixture was homogenised for 20 minutes.

The size, the bulked and tapped density, Carr Index and Flow Index of the resulting granules were determined according to the aforementioned procedures. The results are given in Table 1:

5     Example 2: Preparation of Feed Supplement Containing the iso propyl ester of 2-hydroxy-4-(methylthio) butanoic acid (HMBI Supported on Sepiolite Clay)

A weight of 27kg of sepiolite clay (70 weight %) was placed in a 130 litres Lodgie plow mixer. The clay was stirred for five minutes. 11.6kg of the liquid isopropyl ester of 2-hydroxy-4-(methylthio) butanoic acid (30weight %) was  
10     poured into the mixer at a flow rate of 36kg/hour. The resulting mixture was homogenised for 20 minutes.

The size, the bulked and tapped density, Carr Index and Flow Index of the resulting granules were determined according to the aforementioned procedures. The results are given in Table 2:

15     Example 3: Preparation of Feed Supplement Containing the iso propyl ester of 2-hydroxy-4-(methylthio) butanoic acid (HMBI Supported on Sepiolite Clay)

A weight of 0.22kg of sepiolite clay (70 weight %) was placed in an Aeromatic fluidized bed. Gas was allowed to flow through the bed. The clay was  
20     suspended in the gas flow for 5 minutes. 0.11kg of liquid isopropyl ester of 2-hydroxy-4-(methylthio) butanoic acid (30weight %) was sprayed through a nozzle positioned at the top of the bed. The spraying procedure was allowed to continue for 10 minutes. Fluidisation of the bed was allowed to continue for a further five minutes thereafter. The gas flow was then discontinued.

25     320g of powdered product was collected and divided by a Retsch device into 8 portions of 40g. Each portion was then divided into 8 further portions of 5g for stability measurement

The stability of the solid product was determined by measuring the amount of hmbi in the solid product after production (0 days) and after 15, 45 and 90 days  
30     The solid material was kept under ambient conditions throughout the test period.

The level of hmbi remained constant throughout the test period indicating good stability of the feed supplement. The results are given in Table 3.

Table 1

Characteristic	
HMB expected theoretical amount	26.4
HMB amount measured	27.4
Size distribution	sieve
D <sub>10</sub> (μm)	150
D <sub>50</sub> (μm)	340
D <sub>90</sub> (μm)	530
% > 630 μm	1.8
% < 100 μm	0.3
Porosity	
Total porous volume (cm <sup>3</sup> /g)	0.51
Residual intraparticle porous volume (cm <sup>3</sup> /g)	0.18
Diameter of interparticle pores (μm)	93
Calculated particle density (g/cm <sup>3</sup> )	1.44
Bulk powder density (g/cm <sup>3</sup> )	0.84
Tapped powder density (g/cm <sup>3</sup> )	0.94
Carr index	12%
Flowing index (annular cell)	23

5

10

15



Table 2

Characteristic	Batch 1	Batch 2
<b>HMBI expected theoretical amount</b>	27.6	27.6
<b>HMBI amount measured par dosage HPLC</b>	28.4	26.3
<b>Size distribution</b>	<b>Sieve</b>	<b>Sieve</b>
D <sub>10</sub> (μm)	167	164
D <sub>50</sub> (μm)	344	356
D <sub>90</sub> (μm)	529	563
% > 630 μm	1.6	2.1
% < 100 μm	1.9	1.7
<b>Porosity</b>		
Total porous volume (cm <sup>3</sup> /g)	0.38	0.38
Residual intraparticle porous volume (cm <sup>3</sup> /g)	0.09	0.10
Powder porosity (%)	42%	42%
Diameter of interparticle pores (μm)	67	73
<b>Calculated particle density (g/cm<sup>3</sup>)</b>	1.44	1.42
<b>Bulk powder density (g/cm<sup>3</sup>)</b>	0.89	0.87
<b>Tapped powder density (g/cm<sup>3</sup>)</b>	0.95	0.93
<b>Carr index</b>	7%	7%
<b>Flowing index ( annular cell)</b>	20	19

Table 3

Amount	Batch
HMBI (theoretical)	30.7%
HMBI (0 days)	31.0 %
HMBI (15 days)	29.8%
HMBI (45 days)	31.2%
HMBI (90 days)	31.4%

### Bio-availability Tests

5

The solid feed supplement containing HMBI prepared according to Example 3 above comprising 30% HMBI and 70% sepiolite clay was added to a feed ration and given to 4 dry cows over a period of 7 days at two different daily doses. A similar study was carried out using liquid HMBI (Comparative Example 1) and the commercially known product Smartamine (Solid form - Comparative Example 2). The amount of feed supplement in each case was calculated to provide 50g of either methionine-equivalent (HMB), in the case of the solid and liquid powdered HMBI or methionin ein the case of Smartamine.

10

The feed supplement was introduced into the normal feed ration once per day. Blood samples were taken every half hour for the first two hours following administration of the feed and then every two hours thereafter for the Example 1 and Comparative Example 1 and every two hours in Comparative Example 2

15

The concentration of methionine in the blood was measured and plotted . the area under the curve was calculated to give the bio-availability results. The feed supplement according to the present invention provided 56% bio-available methionine. The liquid hmbi feed supplement provided 49% bio-avaialble methionine. Smartamine provided 82% bio-available methionine. The liquid and solid hmbi feed supplements resulted in methionine absorption into the bloodstream within 2 hours after intake whilst for Smartamine, methionine was detected after 10 hours. resulted in methionine.

20

25

The results are plotted in Figure 1 where:

HMBIP80 is solid HMBI fed at a dose providing 80g methionine-equivalent (or HMB) per cow per day ("P" stands for powder).

5 HMBIP50 is solid HMBI fed at a dose providing 50g methionine-equivalent (HMB) per cow per day

HMBIL is liquid HMBI fed at a dose providing 50g methionine-equivalent (HMB) per cow per day (L = liquid)

Sm.M is Smartamine M, fed at a dose providing 50 g methionine per cow per day.

10

Also:

- HMBI is the abbreviation for "isopropyl ester of 2-hydroxy-4 (methylthio) butanoic acid".
- HMB is the abbreviation for "2-hydroxy-4 (methylthio) butanoic acid".

15

## CLAIMS

5 1. An animal feed supplement in the solid form said feed supplement comprising 2-hydroxy-4-(methylthio) butanoic acid or an ester thereof and a porous carrier material wherein the 2-hydroxy-4-(methylthio) butanoic acid or ester thereof accounts for at least 20% by weight of the total composition of the feed supplement.

10 2. A feed supplement as claimed in claim 1 in which the 2-hydroxy-4-(methylthio) butanoic acid or an ester thereof is present in an amount of from 20 to 40% by weight.

15 3. A feed supplement as claimed in claim 1 or claim 2 in which the ester is a linear or branched alkyl ester selected from methyl ester, ethyl ester, n-propyl ester, isopropyl ester, butyl esters, namely n-butyl ester, sec butyl ester, isobutyl ester and tertiary butyl ester, pentyl esters and hexyl esters, especially n-pentyl, isopentyl, n-hexyl and isohexyl esters.

4. A feed supplement as claimed in claim 3 in which the ester is the isopropyl ester or the tertiary butyl ester

20 5. A feed supplement as claimed in any one of the preceding claims in which the porous carrier material has a porosity of from 0.25 ml/g.

6. A feed supplement as claimed in any one of the preceding claims in which the carrier material is selected from clays such as vermiculite, sepiolite and perlite clays; silicates and silica such as magnesium silicates and zeolites.

25 7. A feed supplement as claimed in claim 6 in which the carrier material is sepiolite.

8. A feed supplement as claimed in any one of the preceding claims in which the particles have a size range of from 200 to 400 microns.


9. An animal feed comprising a feed supplement as claimed in any one of the preceding claims.

## ABSTRACT

## ANIMAL FEED SUPPLEMENT

5     An animal feed supplement in the solid form said feed supplement comprising 2-  
hydroxy-4-(methylthio) butanoic acid or an ester thereof and a porous carrier  
material wherein the 2-hydroxy-4-(methylthio) butanoic acid or ester thereof  
accounts for at least 20% by weight of the total composition of the feed  
supplement.

10



5 Figure 1 Blood plasma methionine concentrations of dairy cows fed various methionine-supplying feed supplements